

## Cleveland State University EngagedScholarship@CSU

Undergraduate Research Posters 2015

Undergraduate Research Posters

2015

# Characterization of the detailed interaction interface between T. brucei telomere proteins TRF and TIF2

Jennifer Kungle  
*Cleveland State University*

John Sabljic  
*Cleveland State University*

Tia Nikova  
*Cleveland State University*

Follow this and additional works at: [https://engagedscholarship.csuohio.edu/u\\_poster\\_2015](https://engagedscholarship.csuohio.edu/u_poster_2015)

**How does access to this work benefit you? Let us know!**

### Recommended Citation

Kungle, Jennifer; Sabljic, John; and Nikova, Tia, "Characterization of the detailed interaction interface between T. brucei telomere proteins TRF and TIF2" (2015). *Undergraduate Research Posters 2015*. 7.  
[https://engagedscholarship.csuohio.edu/u\\_poster\\_2015/7](https://engagedscholarship.csuohio.edu/u_poster_2015/7)

This Book is brought to you for free and open access by the Undergraduate Research Posters at EngagedScholarship@CSU. It has been accepted for inclusion in Undergraduate Research Posters 2015 by an authorized administrator of EngagedScholarship@CSU. For more information, please contact [library.es@csuohio.edu](mailto:library.es@csuohio.edu).



This digital edition was prepared by MSL Academic Endeavors, the imprint of the Michael Schwartz Library at Cleveland State University.

## ***Characterization of the detailed interaction interface between T. brucei telomere proteins TRF and TIF2***

College of Sciences and Health Professions

**Student Researchers:** Jennifer Kungle, John Sabljic, and Tia Nikova

**Faculty Advisor:** Bibo Li

### **Abstract**

The protozoan parasite *Trypanosoma brucei* causes fatal African trypanosomiasis in humans and nagana in cattle. *T. brucei* switches its variant surface glycoproteins (VSGs) inside the mammalian host, evading the host immune response. VSGs are expressed monoallelically from subtelomeric expression sites, and telomere proteins regulate VSGs.

We previously found that telomere protein *TbTIF2* interacts with *TbTRF* (TTAGGG-repeat binding factor) and plays important roles in VSG switching regulation. *TbTRF* maintains the telomere terminal structure. *TbTIF2* is essential for subtelomeric integrity and suppresses VSG switching by inhibiting subtelomeric gene conversion. Depletion of *TbTIF2* decreases *TbTRF* protein level. We hypothesize that *TbTRF-TbTIF2* interaction is essential for maintaining *TbTRF* protein level. We test this hypothesis by mapping the interaction between *TbTRF* and *TbTIF2*. *TbTRF* has both N-terminal TRF Homology (TRFH) and C-terminal Myb domains. *TbTRFH* contains seven helices and interacts with the N-terminus of *TbTIF2* (aa 2-190). We found that deleting either of the first two helices in *TbTRFH* abolishes its interaction with *TbTIF2*. Currently we are generating deletion and point mutations within the first two helices of *TbTRFH*, which will be tested for their ability to interact with *TbTIF2* to determine the key residues in *TbTRFH* that are required for interacting with *TbTIF2*.

*\*Partially Supported by the McNair Scholars Program*